

Molecular Line Lists For Exoplanets and Other Atmospheres

Jonathan Tennyson

*Department of Physics and Astronomy, University College London,
London WC1E 6BT, United Kingdom*

Abstract. At elevated temperatures the spectra of polyatomic molecules become extremely complicated with millions, or even billions, of transitions potentially playing an important role. The atmospheres of cool stars and "hot Jupiter" extrasolar planets are rich with molecules in the temperature range 1000 to 3000 K and their properties are strongly influenced by the infrared and visible spectra of these molecules. So far there are extensive, reliable lists of spectral lines for a number species including some stable diatomics, water, ammonia. However data is almost completely lacking for many key species such as methane.

The ExoMol project (www.exomol.com) aims to construct line lists of molecular transitions suitable for spectroscopic and atmospheric modelling of cool stars and exoplanets [1]. At elevated temperatures it is necessary to consider many millions, even billions, of lines for a single species. Line lists therefore are computed on the basis of robust theoretical models tested against available laboratory data rather than constructed experimentally. Illustrative examples will be discussed and as will progress on obtaining the full set of molecular data. Given the sheer volume of data it is also necessary to consider data representation and storage issues.

ACKNOWLEDGMENTS

This work was performed as part of ERC Advanced Investigator Project 267219 and the project VAMDC which is funded by the European Union INFRA-2008-1.2.2 Scientific Data Infrastructure program under Grant Agreement number 239108.

REFERENCES

1. J. Tennyson and S. N., Yurchenko, *Mon. Not. Roy. Astron. Soc.* 2012, (in press). arXiv:1204.