

Autoionizing Rates for Excited States of Many-Electron Ions

U.I. Safronova and W.R. Johnson

Department of Physics, University of Notre Dame, Notre Dame, IN 46556

1 Introduction

The most important difference between autoionization states and other excited states is an additional broadening of levels caused by the decay of these states (e.g. $2\ell n\ell' \Rightarrow 1sk\ell''$). One can express the autoionizing widths of a level as the square of the decay amplitude:

$$,^N(aLS, a'LS) = \sum_{a_0kl} \gamma(aLS, a_0klLS) \gamma(a'LS, a_0klLS), \quad (1)$$

where

$$\gamma(aLS, a_0klLS) = \sqrt{\frac{2\pi}{k}} \left\langle aLS \left| \sum_{i>j} 1/r_{ij} \right| a_0klLS \right\rangle. \quad (2)$$

Here k is the momentum of the ejected electron in the continuum state (a_0klLS), and aLS describes the autoionizing state. When a nonrelativistic basis is used, relativistic corrections must be taken into account perturbatively. A nonrelativistic basis was used for calculations of autoionization rates in the MZ code [1]–[2], in the model-potential method (SUPERSTRUCTURE code) [3], and in the multiconfiguration Hartree-Fock method (Cowan code [4]).

The purpose of this paper is to obtain simple formulas for autoionization rates for any atomic system by averaging over (LSJ). These calculations will be carried out using screened hydrogenic wave functions. The purpose of these calculations is to provide accurate, but easily used, atomic data.

2 Results and Discussion

Autoionization changes the state of two atomic electrons which initially can belong to either two different groups or to the same group of equivalent electrons. One of the electrons makes a transition to the continuous spectrum, and the other changes its state in the atom. Therefore, to calculate the probability of an autoionizing decay in a general multi-electron ion, it is sufficient to study the change of state in two systems: $(n_1l_1)^{p_1}(n_2l_2)^{p_2}(n_3l_3)^{p_3}$ and $(n_1l_1)^{p_1}(n_2l_2)^{p_2}$. We must study two types of transitions:

$$\begin{aligned} (n_1l_1)^{p_1}(n_2l_2)^{p_2}(n_3l_3)^{p_3} &\Rightarrow (n_1l_1)^{p_1-1}(n_2l_2)^{p_2-1}(n_3l_3)^{p_3+1}k\ell \\ (n_1l_1)^{p_1}(n_2l_2)^{p_2} &\Rightarrow (n_1l_1)^{p_1-2}(n_2l_2)^{p_2+1}k\ell. \end{aligned} \quad (3)$$

For LS averaged autoionization decay probabilities, one obtains in each case:

$$\begin{aligned} ,^N \left[(n_1l_1)^{p_1}(n_2l_2)^{p_2}(n_3l_3)^{p_3} \Rightarrow (n_1l_1)^{p_1-1}(n_2l_2)^{p_2-1}(n_3l_3)^{p_3+1}k\ell \right] = \\ p_1p_2 \left(1 - \frac{p_3}{N_3} \right) A(n_1l_1n_2l_2; n_3l_3k\ell), \end{aligned} \quad (4)$$

and

$$, {}^N \left[(n_1 l_1)^{p_1} (n_2 l_2)^{p_2} \Rightarrow (n_1 l_1)^{p_1-2} (n_2 l_2)^{p_2+1} k \ell \right] = \frac{1}{2} p_1 (p_1 - 1) \left(1 - \frac{p_2}{N_2} \right) A(n_1 l_1 n_1 l_1; n_2 l_2 k \ell), \quad (5)$$

where $N_i = 2(2l_i + 1)$ and where $A(n_1 l_1 n_2 l_2; n_3 l_3 k \ell)$ is the two-electron decay probability.

$$\begin{aligned} & A(n_1 l_1 n_2 l_2; n_3 l_3 k l_4) \\ = & \frac{2\pi}{k} (2l_3 + 1)(2l_4 + 1) \sum_l \left[\frac{1}{(2l + 1)} P_l^2(n_1 l_1 n_2 l_2; n_3 l_3 k l_4) + \frac{1}{(2l + 1)} P_l^2(n_1 l_1 n_2 l_2; k l_4 n_3 l_3) \right. \\ & \left. - \sum_{l'} \left\{ \begin{matrix} l_1 & l_4 & l' \\ l_2 & l_3 & l \end{matrix} \right\} P_l(n_1 l_1 n_2 l_2; n_3 l_3 k l_4) P_{l'}(n_1 l_1 n_2 l_2; k l_4 n_3 l_3) \right] \end{aligned} \quad (6)$$

$$\begin{aligned} A(n_1 l_1 n_1 l_1; n_3 l_3 k l_4) = & \frac{2\pi}{k} \frac{(4l_1 + 1)}{(2l_1 + 1)} (2l_3 + 1)(2l_4 + 1) \sum_l \left[\frac{2}{(2l + 1)} P_l^2(n_1 l_1 n_2 l_2; n_3 l_3 k l_4) \right. \\ & \left. - \sum_{l'} \left\{ \begin{matrix} l_1 & l_4 & l' \\ l_1 & l_3 & l \end{matrix} \right\} P_l(n_1 l_1 n_1 l_1; n_3 l_3 k l_4) P_{l'}(n_1 l_1 n_1 l_1; k l_4 n_3 l_3) \right] \end{aligned} \quad (7)$$

Here P_l can be described as a product of a radial integral and 3j-Wigner coefficients:

$$P_l(n_1 l_1 n_1 l_1; n_3 l_3 k l_4) = R_l(n_1 l_1 n_1 l_1; n_3 l_3 k l_4) (-1)^{l+(l_1+l_2+l_3+l_4)/2} \begin{pmatrix} l_1 & l_4 & l \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_2 & l_3 & l \\ 0 & 0 & 0 \end{pmatrix} \quad (8)$$

Results of our calculations of $A(n_1 l_1 n_2 l_2; n_3 l_3 k l_4)$ and $A(n_1 l_1 n_1 l_2; n_3 l_3 k l_4)$ are given in Tables 1 and 2 for $n_1 l_1, n_2 l_2$ with $7 \geq n_1 \geq 1, (n_1 - 1) \geq l_1 \geq 0, 7 \geq n_2 \geq 1, (n_2 - 1) \geq l_2 \geq 0$. The value of n_3 was chosen such that $\sqrt{\frac{1}{n_3^2} - \frac{1}{n_1^2} - \frac{1}{n_2^2}} > 0$ and $(n_3 - 1) \geq l_3 \geq 0$. The value of l_4 was defined by the triangle rule for the 3j-Wigner coefficient in Eq. (9). The value of k was given by $k = \frac{Z}{Z-\sigma} \sqrt{\frac{1}{n_3^2} - \frac{1}{n_1^2} - \frac{1}{n_2^2}}$, where Z is the nuclear charge and σ is a screening constant.

In total, 8116 kinds of $A(n_1 l_1 n_2 l_2; n_3 l_3 k l_4)$ and $A(n_1 l_1 n_1 l_2; n_3 l_3 k l_4)$ were calculated. We have selected the results with the largest values of A for illustration.

Acknowledgements

The work was supported by Grant No. B336454 from the Lawrence Livermore National Laboratory.

References

- [1] L. A. Vainshtein and U. I. Safronova, *At. Data Nucl. Data Tables* **21**, 49 (1978); *ibid.* **25**, 311 (1980)
- [2] U. I. Safronova, *Physica Scripta* **T26**, 59 (1989)
- [3] F. Bely-Dubau, A. N. Gabriel and S. Volonte, *Mon. Not. R. Astron. Soc.* **186**, 405 (1979), *ibid.* **189**, 801 (1979)
- [4] R. D. Cowan, *The Theory of Atomic Structure and Spectra* (Berkeley, 1981)

Table 1: Autoionizing rates $A(n_1l_1n_2l_2; n_3l_3kl_4)$ in s^{-1} as function of Z and σ

$n_1l_1n_2l_2$	$n_3l_3kl_4$	$Z = 26$ $\sigma = 0$	$Z = 26$ $\sigma = 1$	$Z = 16$ $\sigma = 1$	$Z = 10$ $\sigma = 1$	$Z = 6$ $\sigma = 1$	$Z = 6$ $\sigma = 2$
$2s2s$	$1sks$	2.057E+14	2.004E+14	1.968E+14	1.908E+14	1.785E+14	1.364E+14
$2s2p$	$1skp$	6.088E+13	5.886E+13	5.753E+13	5.533E+13	5.098E+13	3.691E+13
$2p2p$	$1sks$	1.039E+13	9.544E+12	9.014E+12	8.184E+12	6.716E+12	3.316E+12
$2p2p$	$1skd$	1.245E+14	1.160E+14	1.104E+14	1.016E+14	8.548E+13	4.511E+13
$2s3s$	$1sks$	2.604E+13	2.528E+13	2.477E+13	2.390E+13	2.212E+13	1.613E+13
$2s3p$	$1skp$	1.736E+13	1.667E+13	1.621E+13	1.545E+13	1.396E+13	9.349E+12
$2p3s$	$1skp$	1.252E+13	1.220E+13	1.199E+13	1.162E+13	1.084E+13	7.853E+12
$2p3p$	$1skd$	2.381E+13	2.208E+13	2.097E+13	1.918E+13	1.591E+13	7.845E+12
$2s4s$	$1sks$	1.050E+13	1.018E+13	9.964E+12	9.599E+12	8.852E+12	6.346E+12
$3s3s$	$1sks$	1.284E+13	1.245E+13	1.218E+13	1.172E+13	1.078E+13	7.601E+12
$3p3p$	$2pkp$	2.805E+14	2.776E+14	2.757E+14	2.723E+14	2.654E+14	2.394E+14
$3p3d$	$2pkd$	1.576E+14	1.565E+14	1.558E+14	1.545E+14	1.517E+14	1.402E+14
$3d3d$	$2pkf$	4.147E+14	4.057E+14	3.997E+14	3.895E+14	3.688E+14	2.981E+14
$3s3s$	$2pkp$	3.169E+13	3.106E+13	3.063E+13	2.988E+13	2.828E+13	2.212E+13
$3s3p$	$2pks$	8.273E+13	8.207E+13	8.162E+13	8.084E+13	7.922E+13	7.306E+13
$3s3d$	$2pkf$	1.811E+13	1.823E+13	1.831E+13	1.846E+13	1.878E+13	2.019E+13
$3p3d$	$2pks$	1.839E+13	1.791E+13	1.758E+13	1.704E+13	1.595E+13	1.239E+13
$3p3d$	$2pkg$	1.138E+13	1.130E+13	1.125E+13	1.115E+13	1.090E+13	9.623E+12
$3d3d$	$2pkp$	3.333E+13	3.251E+13	3.196E+13	3.104E+13	2.918E+13	2.303E+13
$3s4p$	$2pks$	1.794E+13	1.768E+13	1.750E+13	1.719E+13	1.655E+13	1.411E+13
$3p4s$	$2pks$	3.325E+13	3.261E+13	3.216E+13	3.141E+13	2.985E+13	2.418E+13
$3p4p$	$2pkp$	7.195E+13	7.028E+13	6.915E+13	6.723E+13	6.328E+13	4.941E+13
$3p4d$	$2pkd$	5.530E+13	5.411E+13	5.330E+13	5.190E+13	4.896E+13	3.810E+13
$3d4p$	$2pkd$	4.024E+13	4.021E+13	4.015E+13	3.994E+13	3.919E+13	3.367E+13
$3d4d$	$2pkp$	1.042E+13	9.845E+12	9.470E+12	8.863E+12	7.716E+12	4.557E+12
$3d4d$	$2pkf$	9.831E+13	9.423E+13	9.150E+13	8.692E+13	7.781E+13	4.963E+13
$3d4f$	$2pkg$	4.169E+13	3.854E+13	3.653E+13	3.336E+13	2.765E+13	1.390E+13
$3p5s$	$2pks$	1.665E+13	1.625E+13	1.598E+13	1.552E+13	1.456E+13	1.114E+13
$3p5p$	$2pkp$	3.318E+13	3.225E+13	3.161E+13	3.054E+13	2.835E+13	2.081E+13
$3p5d$	$2pkd$	2.643E+13	2.570E+13	2.520E+13	2.434E+13	2.256E+13	1.627E+13
$3d5p$	$2pkd$	1.731E+13	1.742E+13	1.746E+13	1.746E+13	1.719E+13	1.422E+13
$3d5d$	$2pkf$	4.263E+13	4.067E+13	3.934E+13	3.711E+13	3.264E+13	1.909E+13
$3d5f$	$2pkg$	2.425E+13	2.221E+13	2.091E+13	1.887E+13	1.524E+13	6.866E+12
$3p6p$	$2pkp$	1.828E+13	1.772E+13	1.734E+13	1.669E+13	1.538E+13	1.092E+13
$3p6d$	$2pkd$	1.479E+13	1.433E+13	1.402E+13	1.349E+13	1.239E+13	8.578E+12
$3d6d$	$2pkf$	2.276E+13	2.168E+13	2.094E+13	1.969E+13	1.718E+13	9.609E+12
$3d6f$	$2pkg$	1.450E+13	1.323E+13	1.242E+13	1.115E+13	8.892E+12	3.793E+12
$3p7p$	$2pkp$	1.120E+13	1.084E+13	1.059E+13	1.018E+13	9.340E+12	6.502E+12
$3d7d$	$2pkf$	1.369E+13	1.303E+13	1.258E+13	1.181E+13	1.025E+13	5.587E+12
$4p4p$	$2pkp$	2.440E+13	2.366E+13	2.316E+13	2.231E+13	2.057E+13	1.454E+13
$4p4d$	$2pkd$	1.351E+13	1.332E+13	1.316E+13	1.288E+13	1.218E+13	8.990E+12
$4d4d$	$2pkf$	2.875E+13	2.722E+13	2.619E+13	2.449E+13	2.116E+13	1.153E+13
$4d5d$	$2pkf$	1.117E+13	1.054E+13	1.012E+13	9.419E+12	8.040E+12	4.119E+12

Table 2: Autoionizing rates $A(n_1l_1n_2l_2; n_3l_3kl_4)$ in s^{-1} as function of Z and σ

$n_1l_1n_2l_2$	$n_3l_3kl_4$	$Z = 26$ $\sigma = 0$	$Z = 26$ $\sigma = 1$	$Z = 16$ $\sigma = 1$	$Z = 10$ $\sigma = 1$	$Z = 6$ $\sigma = 1$	$Z = 6$ $\sigma = 2$
4f5f	3dkg	1.647E+14	1.626E+14	1.612E+14	1.587E+14	1.534E+14	1.338E+14
4s5d	3dks	1.572E+13	1.565E+13	1.560E+13	1.551E+13	1.533E+13	1.464E+13
4p5d	3dkp	2.090E+13	2.081E+13	2.075E+13	2.064E+13	2.040E+13	1.947E+13
4d5s	3dks	3.719E+13	3.697E+13	3.682E+13	3.656E+13	3.600E+13	3.385E+13
4d5p	3dkp	4.649E+13	4.619E+13	4.599E+13	4.564E+13	4.491E+13	4.205E+13
4d5d	3dkd	8.276E+13	8.211E+13	8.166E+13	8.090E+13	7.928E+13	7.309E+13
4d5f	3dkf	8.039E+13	7.983E+13	7.944E+13	7.878E+13	7.735E+13	7.172E+13
4d5g	3dkg	1.309E+13	1.291E+13	1.279E+13	1.259E+13	1.217E+13	1.076E+13
4f5p	3dks	1.263E+13	1.232E+13	1.212E+13	1.178E+13	1.108E+13	8.755E+12
4f5p	3dkd	7.060E+12	7.362E+12	7.569E+12	7.922E+12	8.659E+12	1.133E+13
4f5d	3dkp	1.789E+13	1.752E+13	1.728E+13	1.686E+13	1.602E+13	1.312E+13
4f5d	3dkf	6.984E+13	6.990E+13	6.992E+13	6.993E+13	6.987E+13	6.858E+13
4f5f	3dkd	1.575E+13	1.548E+13	1.529E+13	1.497E+13	1.432E+13	1.203E+13
4f5g	3dkh	9.714E+13	9.476E+13	9.317E+13	9.049E+13	8.509E+13	6.713E+13
4d6s	3dks	2.084E+13	2.057E+13	2.037E+13	2.005E+13	1.935E+13	1.669E+13
4d6p	3dkp	2.589E+13	2.552E+13	2.527E+13	2.483E+13	2.391E+13	2.038E+13
4d6d	3dkd	4.157E+13	4.085E+13	4.035E+13	3.951E+13	3.773E+13	3.111E+13
4d6f	3dkf	4.038E+13	3.970E+13	3.923E+13	3.842E+13	3.671E+13	3.020E+13
4f6d	3dkp	1.066E+13	1.017E+13	9.847E+12	9.316E+12	8.292E+12	5.290E+12
4f6d	3dkf	3.347E+13	3.368E+13	3.379E+13	3.391E+13	3.391E+13	3.146E+13
4f6f	3dkg	7.623E+13	7.435E+13	7.305E+13	7.080E+13	6.607E+13	4.889E+13
4f6g	3dkh	5.976E+13	5.693E+13	5.505E+13	5.197E+13	4.600E+13	2.856E+13
4f6h	3dkk	1.726E+13	1.595E+13	1.512E+13	1.380E+13	1.142E+13	5.689E+12
4d7s	3dks	1.288E+13	1.265E+13	1.250E+13	1.224E+13	1.168E+13	9.556E+12
4d7p	3dkp	1.594E+13	1.565E+13	1.545E+13	1.510E+13	1.436E+13	1.156E+13
4d7d	3dkd	2.443E+13	2.388E+13	2.351E+13	2.286E+13	2.152E+13	1.660E+13
4d7f	3dkf	2.368E+13	2.314E+13	2.278E+13	2.215E+13	2.082E+13	1.591E+13
4f7d	3dkf	1.914E+13	1.936E+13	1.947E+13	1.959E+13	1.958E+13	1.733E+13
4f7f	3dkg	4.287E+13	4.162E+13	4.075E+13	3.922E+13	3.596E+13	2.430E+13
4f7g	3dkh	3.744E+13	3.531E+13	3.390E+13	3.160E+13	2.718E+13	1.495E+13
4f7h	3dkk	1.526E+13	1.390E+13	1.304E+13	1.169E+13	9.325E+12	4.040E+12
5d5d	3dkd	2.636E+13	2.581E+13	2.543E+13	2.478E+13	2.342E+13	1.828E+13
5d5f	3dkf	2.233E+13	2.212E+13	2.196E+13	2.166E+13	2.091E+13	1.717E+13
5f5f	3dkg	4.754E+13	4.584E+13	4.468E+13	4.273E+13	3.876E+13	2.564E+13
5f5g	3dkh	1.310E+13	1.246E+13	1.204E+13	1.135E+13	1.004E+13	6.215E+12
5d6d	3dkd	1.117E+13	1.087E+13	1.067E+13	1.033E+13	9.598E+12	6.887E+12
5d6f	3dkf	1.066E+13	1.049E+13	1.036E+13	1.012E+13	9.566E+12	7.094E+12
5f6d	3dkf	1.035E+13	1.029E+13	1.024E+13	1.010E+13	9.723E+12	7.484E+12
5f6f	3dkg	1.995E+13	1.909E+13	1.851E+13	1.752E+13	1.552E+13	9.117E+12
5f7f	3dkg	1.109E+13	1.058E+13	1.024E+13	9.646E+12	8.439E+12	4.635E+12
5f7s	4fks	2.128E+13	2.122E+13	2.118E+13	2.112E+13	2.097E+13	2.041E+13
5f7p	4fkp	2.444E+13	2.437E+13	2.433E+13	2.425E+13	2.408E+13	2.341E+13
5f7d	4fkf	3.005E+13	2.995E+13	2.989E+13	2.977E+13	2.953E+13	2.857E+13
5f7f	4fkf	4.241E+13	4.225E+13	4.214E+13	4.195E+13	4.154E+13	3.993E+13