

## **A Study of tungsten spectra using Large Helical Device and Compact Electron Beam Ion Trap in NIFS**

S.Morita<sup>a,b</sup>, C.F.Dong<sup>a</sup>, M.Goto<sup>a,b</sup>, M.Hasuo<sup>c</sup>, D.Kato<sup>a</sup>, F.Koike<sup>d</sup>, I.Murakami<sup>a</sup>,  
N.Nakamura, T.Oishi<sup>a</sup>, H.A.Sakaue<sup>a</sup>, A.Sasaki<sup>f</sup>, C.Suzuki<sup>a</sup> and E.H.Wang<sup>b</sup>

<sup>a</sup>National Institute for Fusion Science, Toki 509-5292, Gifu, Japan

<sup>b</sup>Department of Fusion Science, Graduate University for Advanced Studies, Toki  
509-5292, Gifu, Japan

<sup>c</sup>Department of Mechanical Engineering and Science, Graduate School of Engineering,  
Kyoto University, Kyoto 606-8501, Japan

<sup>d</sup>Physics Laboratory, School of Medicine, Kitasato University, Sagamihara 252-0374,  
Japan

<sup>e</sup>Institute of Laser Science, University of Electro-Communications, Tokyo 182-8585,  
Japan

<sup>f</sup>Quantum Beam Science Directorate, Japan Atomic Energy Research Agency, Kizugawa  
619-0215, Kyoto, Japan

E-mail address: morita@nifs.ac.jp

Tungsten spectra have been observed from Large Helical Device (LHD) and Compact electron Beam Ion Trap (CoBIT) in wavelength range of visible to EUV. An impurity pellet (cylindrical carbon with tungsten) is injected to LHD plasmas for observing the tungsten spectra. Radial profiles of EUV spectra from highly ionized tungsten ions have been measured and analyzed with impurity transport simulation code to examine the ionization balance determined by ionization and recombination rate coefficients. In order to observe the tungsten spectra from lower-ionized stages, which can give useful information on the tungsten influx in fusion plasmas, the ablation cloud of the impurity pellet is directly measured with visible spectroscopy. A lot of spectra from neutral and singly ionized tungsten are observed and some of them are identified. A magnetic forbidden line from highly ionized ions of high-Z elements including tungsten observed from LHD plasmas has been analyzed for reconstruction of the atomic structure model in addition to the diagnostic use for D-T burning plasmas. The EUV spectra of tungsten ions, e.g.,  $W^{+24-26}$ , measured from LHD plasmas are compared with those measured from plasmas of CoBIT with monoenergetic electron beam ( $\leq 2\text{keV}$ ). The data are analyzed with collisional-radiative model.